

On the number of connected components of complements to arrangements of subtori

I. N. Shnurnikov*

Abstract

We consider the arrangements of subtori in a flat d -dimensional torus T^d . Let us consider an arrangement on n subtori of codimension one, let f be the number of connected components of the complement in T^d to the union of subtori. We found the set of all possible numbers f for given n and d and arbitrary arrangements of subtori.

MSC: 52C35

Keywords: arrangements of subtori, number of connected components.

Introduction

The theory of plane arrangements in affine or projective spaces has been investigated thoroughly, see the book of Orlic and Terao, [4] and Vassiliev's review [8]. Inspired by the conjecture of Grünbaum [2], Martinov [3] found all possible pairs (n, f) such that there is a real projective plane arrangement of n pseudolines and f regions. It turns out, that some facts concerning the arrangements of hyperplanes or oriented matroids could be generalized to arrangements of submanifolds, see Deshpande [1].

We study the sets $F(T^d, n)$ of connected components numbers of the complements in a flat d -dimensional torus T^d to the unions of n closed connected codimensional one subtori. Author in [7] found the sets $F(T^2, n)$ of region numbers in arrangements of n closed geodesics in a two dimensional torus and a Klein bottle with locally flat metrics. The main result of the present paper is the following:

$$F(T^d, n) = \{l \in \mathbb{N} \mid n - d + 1 \leq l \leq n \text{ or } 2(n - d) \leq l\}$$

for $n > d \geq 2$. Also $F(T^d, n) = \mathbb{N}$ for $2 \leq n \leq d$. $F(T^d, 1) = \{1\}$.

Main part.

Let $\mathbb{R}^d$ be the real d -dimensional vector space and $u_1, \dots, u_d$ are linear independent vectors in $\mathbb{R}^d$. Let L be the set of all linear combinations $k_1 u_1 + \dots + k_d u_d$ with integer coefficients $k_1, \dots, k_d$. The set L could be considered as the group of transitions, acting on $\mathbb{R}^d$. The quotient space of this action is torus, the induced metric (from euclidean metric in $\mathbb{R}^n$) is locally flat. So we call this quotient space *flat torus* T^d and denote by $\mathbb{R}^d/L$. So the factorisation map $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}^d/L$ sends point x to equivalent class $x + L$. Let $(x_1, \dots, x_d)$ are coordinates of $\mathbb{R}^d$ according to the basis u_i . Let us consider a hyperplane K in $\mathbb{R}^d$ with equation $\sum a_i x_i = c$. The subset $\varphi(K) \subset T^d$ is closed iff the numbers $\frac{a_i}{a_j}$ are rational for all pairs i, j such that $a_j \neq 0$. In the set L one may choose another basis $v_1, \dots, v_d$, then the matrix of basis transformation is a $SL(\mathbb{Z})$ matrix.

Let A be the union of n codimensional one subtori in the flat d -dimensional torus T^d . Consider the connected components of the complement $T^d \setminus A$; denote the number of connected components by $f = |\pi_0(T^d \setminus A)|$; let $F(T^d, n)$ be the set of all possible numbers f .

*NRU HSE, Moscow

Lemma 1. Let K be a hyperplane in $\mathbb{R}^d$ with equation $\sum a_i x_i = c$ in a basis $U = (u_i)$ of the lattice for integers a_i such that $\gcd(a_1, \dots, a_d) = 1$.

(a) There is a new basis $V = (v_i)$ of the lattice, according to which K is given by equation $y_1 = 0$. The matrix M , $V = UM$ of basis transfer is a $SL(\mathbb{Z})$ matrix with integer coefficients and the determinant is 1.

(b) Let L be such that U is an orthogonal basis and $|u_i| = 1$ for all i . Then the distance between K and the nearest point $x \in L$ of the lattice such that $x \notin K$ is $\frac{1}{\sqrt{\sum_{i=1}^d a_i^2}}$.

(c) Let U be an orthogonal basis and $|u_i| = 1$ for all i . The volume of T^d is 1. The subset $\varphi(K)$ is a subtorus in T^d and the $(d-1)$ -dimensional volume of $\varphi(K)$ is $\sqrt{\sum_{i=1}^d a_i^2}$.

Lemma 2. Let $A_1, \dots, A_n$ be the set of n codimensional one subtori in the flat d -dimensional torus T^d . Let m be the maximal number of parallel tori among $A_1, \dots, A_n$. Then $f \geq m(n - m - d + 2)$.

Proof. Induction on d , base $d = 1$ is trivial. Suppose the statement is true for $d - 1$ and every n . Let us prove the statement for d . Consider m parallel subtori, they divide torus into m components U_i , $i = 1, \dots, m$ and each component is homeomorphic to prime product of a segment and $(d - 1)$ -dimensional torus. For each U_i let us consider a control $(d - 1)$ -dimensional torus R_i such that the other $n - d$ tori of $A_1, \dots, A_n$ form on R_i an arrangement of $n - m$ subtori and so by the induction assumption they divide R_i into at least $n - m - d + 2$ components. So, each U_i is divided by the other $n - d$ tori of $A_1, \dots, A_n$ into at least $n - m - d + 2$ regions and the induction is over. $\square$

Lemma 3. Let $n > d \geq 2$. Then $f \geq 2n - 2d$ or: $f \leq n$ and there are at least $n - d + 1$ parallel tori.

Proof. Induction on d , base $d = 2$ is found in [7]. Suppose the statement is true for $d - 1$, let us prove it for d . Let m be the maximal number of parallel tori in the arrangement $A = \{A_1, \dots, A_n\}$. By lemma 2 we have $f \geq m(n - m - d + 2)$. So we over in case $2 \leq m \leq n - d$, because in this case $m(n - m - d + 2) \geq 2(n - d)$.

Let us consider the case $m = 1$. Then we consider a control torus R_1 which is parallel to torus A_1 and is in general position to other tori of arrangement. Let us consider arrangement $B = \{B_1, \dots, B_{n-1}\}$, which is formed in R_1 by other tori of arrangement A . We know that $f(A) \geq f(B)$. If in B there are at most $n - d$ parallel tori then by induction assumption we have $f(B) \geq 2(n - d)$. Otherwise in B there is at least $n - d + 1$ parallel tori and this means that there exist a $(d - 2)$ -dimensional torus S^{d-2} which is parallel to at least $n - d + 2$ tori of A . Consider the two-dimensional subtori W^2 which is orthogonal to S^{d-2} . Then by induction assumption for $d = 2$ we have that A divides W^2 into $f(W^2) \geq 2(n - d)$ regions and so $f(A) \geq f(W^2) \geq 2(n - d)$.

Let us consider the case $m \geq n - d + 1$. The subset of m parallel tori divide T^d into m regions. If at least one of them is divided into more then one connected components by the other tori of A , then every of m regions is divided in at least two connected components and so $f(A) \geq 2m \geq 2(n - d)$. In the other case the induction is over. $\square$

Theorem 1.

$$F(T^d, n) = \{l \in \mathbb{N} \mid n - d + 1 \leq l \leq n \text{ or } 2(n - d) \leq l\}$$

for $n > d \geq 2$. Also $F(T^d, n) = \mathbb{N}$ for $2 \leq n \leq d$. $F(T^d, 1) = \{1\}$.

Proof. The case $d = 2$ is proved in [7]. Now we assume $d \geq 3$. From the lemma 3, lemma 2 we see that the set $F(T^d, n)$ does not contain other integers.

We construct examples for $\leq n$ and $\geq 2n - 2d$ regions separately.

Let us consider n hyperplanes in $\mathbb{R}^d$ (an equation corresponds to a hyperplane):

$$\begin{aligned} x_i &= 0, & 1 \leq i \leq k, \\ x_{k+1} &= c_{i-k}, & k + 1 \leq i \leq n \end{aligned}$$

for some integer k , $0 \leq k \leq d-1$ and real c_{i-k} with different fractional parts. By the factorization map $\mathbb{R}^d \rightarrow \mathbb{R}^d/Z^d$ we get a set $\{T_i^{d-1}, i = 1, \dots, n\}$ of n codimensional one subtori. And the complement is homeomorphic to the prime product

$$T^d \setminus \bigcup_i T_i^{d-1} \approx \mathbb{R}^k \times (S^1 \setminus \{p_1, \dots, p_{n-k}\}) \times (S^1)^{d-k-1},$$

where $S^1 \setminus \{p_1, \dots, p_{n-k}\}$ denotes a circle without $n-k$ points. Hence the number of complement regions equals $n-k$, for an integer k such that $0 \leq k \leq d-1$.

Now let us take a nonnegative integer k and construct an arrangement with $2n-2d+k$ connected components of the complement. We determine the subtori by equations:

$$\begin{aligned} x_i &= 0, \quad \text{for } 2 \leq i \leq d, \\ x_2 &= kx_1 + \frac{1}{2}, \\ x_1 &= c_j \quad \text{for } j = 1, \dots, n-d, \end{aligned}$$

whereas numbers $kc_j + \frac{1}{2}$ are not integer for any j . (This means that the intersection of three subtori

$$x_2 = kx_1 + \frac{1}{2}, \quad x_1 = c_j, \quad x_2 = 0$$

is an empty set.) Therefore

$$T^d \setminus \bigcup_{i=3}^d \{x_i = 0\} \approx T^2 \times \mathbb{R}^{d-2}.$$

In the 2-dimensional torus the equations

$$\begin{aligned} x_2 &= 0, \\ x_2 &= kx_1 + \frac{1}{2}, \\ x_1 &= c_j \quad \text{for } j = 1, \dots, n-d \end{aligned}$$

produce the arrangement of $n-d+2$ closed geodesics. The geodesic union divides the torus into $2n-2d+k$ connected components (for more details on the arrangements of closed geodesics in a flat torus see [7]). $\square$

Lemma 4. *Let K_1 and K_2 be hyperplanes in $\mathbb{R}^d$ given by the equations $x_1 = 0$ and $\sum_i a_i x_i = c$ where a_i are integers. Then the intersection of subtori $\varphi(K_1)$ and $\varphi(K_2)$ in T^d consists of $\gcd(a_2, \dots, a_d)$ connected components, each of which is a $(d-2)$ -dimensional subtorus.*

Lemma 5. *Let $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ be integer vectors such that $\gcd(a_1, \dots, a_n) = 1$ and $\gcd(b_1, \dots, b_n) = 1$. Let two subtori in a flat d -dimensional torus are given by vectors $\mathbf{a}$ and $\mathbf{b}$. Let $f(\mathbf{a}, \mathbf{b})$ be the number of connected components of the intersection of two subtori. Then*

$$\begin{aligned} f(\mathbf{a}, \mathbf{b}) = \gcd \left(\frac{a_2 b_1 - b_2 a_1}{\gcd(a_1, a_2)}, \frac{a_3 (b_1 u_1^{(1)} + b_2 u_2^{(1)}) - b_3 \gcd(a_1, a_2)}{\gcd(a_1, a_2, a_3)}, \right. \\ \dots \\ \left. \frac{a_n (b_1 u_1^{(n-2)} + b_2 u_2^{(n-2)} + \dots + b_{n-1} u_{n-1}^{(n-2)}) - b_n \gcd(a_1, \dots, a_{n-1})}{\gcd(a_1, \dots, a_n)} \right) \end{aligned}$$

where and u_i^j are integers such that

$$\gcd(a_1, \dots, a_{j+1}) = a_1 u_1^{(j)} + a_2 u_2^{(j)} + \dots + a_{j+1} u_{j+1}^{(j)}$$

for each $j = 1, \dots, n-2$.

References

- [1] P. Deshpande, Arrangements of Submanifolds and the Tangent Bundle Complement. *Electronic Thesis and Dissertation Repository*, Paper 154 (2011).
- [2] B. Grünbaum, *Arrangements and Spreads*. AMS, Providence, Rhode Island, 1972.
- [3] N. Martinov, Classification of arrangements by the number of their cells. *Discrete and Comput. Geometry* (1993) **9**, N 1, 39–46.
- [4] P. Orlic, H. Terao, Arrangements of Hyperplanes. Springer – Verlag, Berlin – Heidelberg, 1992. 329 pp.
- [5] R.W. Shannon, A lower bound on the number of cells in arrangements of hyperplanes. *Jour. of combinatorial theory (A)*, **20**, (1976) 327–335.
- [6] I. N. Shnurnikov, Into how many regions do n lines divide the plane if at most $n - k$ of them are concurrent? *Moscow Univ. Math. Bull., ser. 1* (2010) **65**:5, 208 – 212.
- [7] I. N. Shnurnikov, On the number of regions formed by arrangements of closed geodesics on flat surfaces, *Math. Notes* **90**, N 3 – 4 (2011), 619 – 622.
- [8] V. A. Vassiliev, Topology of plane arrangements and their complements. *Uspekhi Mat. Nauk*, **56**, iss. 2(338), (2001), 167 — 203.